

Composition:

Timonac™ 50 mg Tablet: Each film coated tablet contains 50 mg of Tiemonium Methylsulphate INN.

Timonac™ 5 mg/2 ml IM/IV Injection: Each 2 ml ampoule contains 5 mg of Tiemonium Methylsulphate INN.

Pharmacology:

Timonac™ is a quaternary ammonium antimuscarinic agent with peripheral effect similar to those of atropine. **Timonac™** strengthens calcium bonding with phospholipids and proteins thus stabilizing the cell membrane of the GI tract.

Indication:

Timonac™ is an antispasmodic drug. It reduces muscle spasms of the intestine, biliary system, uterus & urinary bladder. It is indicated for the pain in gastrointestinal & biliary disease in the urology and gynecology such as gastroenteritis, diarrhea, dysentery, biliary colic, enterocolitis, cholecystitis, colonopathy, mild cystitis, & spasmodic dysmenorrhoea.

Dose & administration:

Route of administration: Oral.

Recommended oral dose of **Timonac™** tablet is 2-6 tablets (100-300 mg) daily in divided dose as required.

Contra-indication:

Tiemonium Methylsulphate tablet should not be used in Glucoma or where acute pain of eyeball with vision disturbance was found. Also this drug should not be used in patients with the disorder of prostate or urinary bladder.

Warning and precaution:

Precautions should be exercised for treatment of patients with disorders of prostate, chronic bronchitis, coronary insufficiency, ambient hyperthermia, cardiac disorders, thyroid disorders, serious renal & hepatic insufficiency.

Side effects:

Side effect with the use of Tiemonium Methylsulphate tablet is very rare.

Use in pregnancy & lactation:

Tiemonium Methylsulphate tablet may be used in pregnancy only if it is clearly needed by the assessment of risk benefit ratio. Tiemonium Methylsulphate tablet may be used in lactating mother only if it is clearly needed by the assessment of risk benefit ratio.

Use in geriatric: Efficacy and safety were maintained with increasing age.

Drug interaction:

Tiemonium Methylsulphate tablet should not be used with other drugs without prior consult of a registered physician to avoid possible drug interactions.

Overdose:

There is no available data regarding the overdose of Tiemonium Methylsulphate.

Storage:

Store in cool & dry place below 30°C, protect from light & moisture. Keep the medicine out of the reach of children.

Packing:

Timonac™ 50 mg Tablet: Each box contains 50 tablets in blister pack.

Timonac™ 5 mg/2 ml IM/IV Injection: Each box contains 10 ampoules in blister pack.

Timonac™ 50 Tablet

Manufactured by:

Jenphar Bangladesh Ltd.

Sreepur, Gazipur, Bangladesh.

Timonac™ IM/IV Injection

Manufactured by:

Jenphar Bangladesh Ltd.

Sreepur, Gazipur, Bangladesh
at Popular Pharmaceuticals Ltd.
Tongi, Gazipur, Bangladesh.